

AD-A074 878

AIR FORCE GEOPHYSICS LAB HANSCOM AFB MA
ATS-5 AND ATS-6 POTENTIALS DURING ECLIPSE. (U)
SEP 79 A G RUBIN, H B GARRETT

F/6 22/1

UNCLASSIFIED

AFOL-TR-79-0218

NL


| OF |
AD
A074878



END
DATE
FILMED

11-79

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 AFGL-TR-79-0218	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 1 2 5
4. TITLE (and Subtitle) 6 <u>ATS-5 AND ATS-6 POTENTIALS DURING ECLIPSE.</u>		5. TYPE OF REPORT & PERIOD COVERED Scientific. Interim.
7. AUTHOR(s) 10 Allen G./Rubin Henry B./Garrett		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Air Force Geophysics Laboratory (PHG) Hanscom AFB Massachusetts 01731 409 578		8. CONTRACT OR GRANT NUMBER(s) LEVEL
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory (PHG) Hanscom AFB Massachusetts 01731		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBER 62-1011 16 76610701 17 077
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 11		12. REPORT DATE 11 27 September 1979
		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 9 Scientific interim rept.,		DDC RECEIVED OCT 11 1979 REGISTERED A
18. SUPPLEMENTARY NOTES Published in NASA Conference Publ. 2071, AFGL-TR-79-0082, 1979		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Spacecraft charging Experimental data ATS-5 ATS-6		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) ATS-5 and ATS-6 data for spacecraft charging during eclipse conditions is analyzed. ATS-5 and ATS-6 charged to voltages greater than 100 volts for about 55% of the eclipse periods examined. The mean spacecraft potential during eclipse was 2 keV for ATS-5, and the highest potential measured was 10 kilovolts. For ATS-6, the mean potential during eclipse was 4 keV, the highest potential measured being 20 keV. The average measured spacecraft potentials for both ATS-5 and 6 depend approximately linearly upon K_p . This		

DD FORM 1473, 1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

409 578

AD A 074878

DDC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

relationship is due mainly to the dependence of electron current density on K_p near midnight. Spacecraft potentials at geosynchronous orbit may, to a rough approximation, thus be inferred from ground-based measurements of K_p , the planetary 3-hour index.

sub p


Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ATS-5 AND ATS-6 POTENTIALS DURING ECLIPSE

Allen G. Rubin and Henry B. Garrett
Air Force Geophysics Laboratory



SUMMARY

ATS-5 and ATS-6 data for spacecraft charging during eclipse conditions is analyzed. ATS-5 and ATS-6 charged to voltages greater than 100 volts for about 55% of the eclipse periods examined. The mean spacecraft potential during eclipse was 2 keV for ATS-5, and the highest potential measured was 10 kilovolts. For ATS-6, the mean potential during eclipse was 4 keV, the highest potential measured being 20 keV. The average measured spacecraft potentials for both ATS-5 and 6 depend approximately linearly upon K_p . This relationship is due mainly to the dependence of electron current density on K_p near midnight. Spacecraft potentials at geosynchronous orbit may, to a rough approximation, thus be inferred from ground-based measurements of K_p , the planetary 3-hour index.

INTRODUCTION

As the ATS-5 and ATS-6 spacecraft have been monitoring the plasma environment at synchronous orbit since 1969, the data provided by these satellites comprises the most extensive data base available of low-energy plasma conditions at geosynchronous orbit.

The present paper is a study of the statistics of charging events based on data from 157 eclipses from ATS-5 and 40 eclipses from ATS-6.

THE DISTRIBUTION OF CHARGING POTENTIALS

For these eclipse charging events, ten minute averages of the charging potentials were obtained. Figure 1 shows the distribution of charging potentials plotted separately for ATS-5 and ATS-6.

For ATS-5, 55% exhibit charging to negative potentials greater than 110 volts and 54% of the ATS-6 charging intervals are greater than 100 volts. This means that both ATS-5 and ATS-6 charge to substantial negative voltages in more than half of the eclipse intervals. The mean charging voltage of ATS-5 is 2 kilovolts for ATS-5 and 4 kilovolts for ATS-6. The highest observed potential is 10 kilovolts for ATS-5 and 20 kilovolts for ATS-6.

Figure 2 shows the distribution of K_p and the 3-hour planetary index

for the ATS-5 and ATS-6 data time periods. ATS-6 data was taken during a more active geomagnetic period than ATS-5, so that the mean value of K_p for the ATS-6 time period is considerably higher than that for the ATS-5 time period. The higher mean charging potential for ATS-6 is no doubt due to hotter injected plasmas for the ATS-6 time period.

CHARGING POTENTIALS VERSUS K_p

Eclipse potentials versus K_p are shown in Figure 3 for eclipse events on ATS-5 and ATS-6 for K_p values up to 6. The error bars shown are the standard deviations ($\pm 1 \sigma$) of the data. The spacecraft potential rises linearly with K_p within the error.

Shown on this figure as well is a theoretical curve based on a current balance calculation in which secondary emission properties of ATS-5 and 6 materials have been accounted for in an approximate manner, using an average secondary emission coefficient. This theoretical curve is given by

$$qV_o = T_e \ln \left(\frac{J_e}{10 J_i} \right)$$

where the factor 10 takes account of secondary emission properties (Garrett and Rubin, 1978).


R.M.S. electron temperature as a function of K_p is shown in Figure 4 for the time period 2100-300. The electron mean temperature varies from 2 to 4 keV. The variation of the ratio of electron to ion current densities with K_p is much greater, as shown in Figure 5. The theoretical prediction of spacecraft potential, as the product of T_e and $\ln (J_e/10 J_i)$, shows that it is the dependence of the ratio of electron to ion current densities on the magnetic activity index, K_p , which is responsible for the strong dependence of charging potential on K_p . The mean R.M.S. electron temperature at a given K_p for the 2100-0300 time period, for the entire ATS-5 and ATS-6 data base, was employed for this calculation. The theoretical curve corresponds very well to observed potentials for K_p values up to 4, but is somewhat higher at larger K_p values.


The spacecraft potential is shown to be proportional to K_p for the 6-hour period around midnight. K_p is an easily accessible quantity, having long been used as an indicator of geomagnetic activity. Since K_p is calculated from ground-based magnetometer systems, it will be useful, to the accuracy of the error bars, as an indicator of the maximum potentials, to be expected on a shadowed spacecraft surface at geosynchronous orbit. As ATS-5 and ATS-6 have radically different shapes, this result could be applicable for a variety of satellites (see Purvis et al., 1977, for a comparison between ATS-5 and ATS-6 potentials in eclipse).


REFERENCES

1. Garrett, H.B. and A.G. Rubin (1978):Spacecraft Charging at Geosynchronous Orbit-Generalized Solution for Eclipse Passage, Geophysical Res. Lett. (to be published).
2. Purvis, C.K., R.O. Bartlett, and S.E. DeForest: Active Control of Spacecraft Charging on ATS-5 and ATS-6, Proceedings of the Spacecraft Charging Technology Conference, edited by C.P. Pike and R.H. Lovell, AFGL-TR-770051/NASA TM 7350, p. 107-120 (1977).


- Figure 1. The occurrence frequency of charging potentials on ATS-5 and ATS-6.
- Figure 2. The occurrence frequency of K_p for ATS-5 and ATS-6 time periods.
- Figure 3. Average eclipse potentials of ATS-5 and ATS-6 versus K_p .
- Figure 4. Mean R.M.S. electron temperature versus K_p for eclipses.
- Figure 5. Ratio of average electron to ion current densities versus K_p for eclipses.







ATS-5
2100-0300



ATS-5
2100-0300

$$\frac{j_e}{10j_i}$$

